


S202 PB


THE OPEN UNIVERSITY
Science: A Second Level Course




Biology

Form and Function

The S202 Picture Book







The S202 Picture Book

The S202 Course Team

This course has been prepared by the following team:

Peggy Varley (*Chairman and General Editor*)

Hendrik Ball (*BBC*)
Gerry Bearman (*Editor*)
Mary Bell
Eric Bowers
Bob Burgoyne
Ian Calvert
Bob Cordell
Norman Cohen
Peter Cole (*BBC*)
Baz East (*Illustrator*)
Vic Finlayson
Anna Furth
Denis Gartside (*BBC*)
Lindsay Haddon
Robin Harding
Stephen Hurry
David Kerrison (*BBC*)

Aileen Llewellyn (*BBC*)
Pam Mullins
Pat Murphy
Seán Murphy
Pam Owen (*Illustrator*)
Phil Parker (*Course Coordinator*)
Ros Porter (*Designer*)
Rob Ransom
Irene Ridge
Steven Rose
Ian Rosenbloom (*BBC*)
Jacqueline Stewart (*Course Editor*)
Mike Stewart
Jeff Thomas
David Tillotson (*Editor*)
Charles Turner
Sue Turner

The Open University Press
Walton Hall, Milton Keynes.

First published 1981. Reprinted 1984, 1988.

Copyright © 1981 The Open University.

All rights reserved. No part of this work may be reproduced in any form, by mimeograph or any other means, without permission in writing from the publisher.

Designed by the Graphic Design Group of the Open University.

Printed in Great Britain by Balding + Mansell UK Limited, London and Wisbech.

ISBN 0 335 16042 5

This text forms part of an Open University course. The complete list of Units in the Course is printed at the end of this text.

For general availability of supporting material referred to in this text please write to: Open University Educational Enterprises Limited, 12 Cofferidge Close, Stony Stratford, Milton Keynes, MK11 1BY, Great Britain.

Further information on Open University courses may be obtained from the Admissions Office, The Open University, P.O. Box 48, Walton Hall, Milton Keynes, MK7 6AB.

contents

How to use this book 2

The Diversity of Organisms

- unit 1 Marine Organisms 3
- unit 2 From Sea to Land: Plants and Arthropods 4
- unit 3 From Sea to Land: Vertebrates 8

Cell Biology

- unit 4 Cell Structure 12
- unit 5 Macromolecules and Membranes 26

Development

- unit 14 Pattern Specification and Morphogenesis 27

Animal Physiology

- unit 16 Communication: Nerves and Hormones 28
- unit 19 Circulatory Systems 32
- units 22 and 23 Osmoregulation and Excretion 33
- units 24 and 25 Nutrition, Feeding Mechanisms and Digestion 34

Plant Physiology

- unit 27 Ion Movement and Phloem Transport 35

Home Experiments

- Animal Histology 37
- Plant Histology 44
- Ferret 45
- Pig's Heart 46
- Cucumbers 47

Index to plates 48

Acknowledgements 50

how to use this book

This book consists of colour pictures and electron micrographs that need to be reproduced by a specialist printer on heavy-grade paper if they are to be seen to best advantage.

The illustrations are grouped according to the subject blocks of the Course and are numbered consecutively. References to the illustrations are generally made in the margins of the Units. We expect you to act on each reference as you come to it, turning first to the relevant pages of this book using the contents list as your guide. If you do not immediately find the picture you need, turn to the index of plates at the back of this book for the 'key words' of the reference, and you will be directed to the plate number (in bold type) and page number of the picture you are seeking.

Note that Unit 4 is different in that it is built around illustrations in this book. It therefore has a different system of reference—see the Study guide to Unit 4.

The *Home Experiment Notes* refer to illustrations directly, by number.

You may of course leaf through *The S202 Picture Book* for general interest and you will also find it provides a helpful back-up to *A Survey of Living Organisms**.

*The Open University (1981) SLO *A Survey of Living Organisms*.




1



2



3




4



5



6



7

1 *Anabaena*, a filamentous blue-green alga, class Cyanophyceae, Kingdom Prokaryota. The larger cells (10 μm) in the middle of the filaments are heterocysts.

2 *Navicula* (60 μm), a diatom, class Bacillariophyceae, division Chrysophyta. The individuals look different because they are viewed from different angles. Markings on the siliceous valves are just visible.


3 *Euglena* (50 μm), division Euglenophyta. Each organism has a long flagellum and many chloroplasts.

4 A solitary blue-green alga (50 μm), class Cyanophyceae, and *Zygnema*, a filamentous green alga, class Chlorophyceae. In *Zygnema*, note the two star-shaped chloroplasts in each cell and the vacuoles; neither of these features is visible in blue-green cells.

5 *Ceratium* (300 μm), a dinoflagellate, division Pyrrophyta, class Dinophyceae. The groove round the middle carries one flagellum and the firm outline is due to the presence of cellulose plates that form three spines.


6 *Chlamydomonas* (20 μm), a unicellular, flagellated green alga, class Chlorophyceae. There is a pair of flagella of equal length and a single chloroplast.

7 *Amoeba* (0.5 mm), phylum Protista, class Sarcodina. Note the single nucleus and spherical contractile vacuole and short pseudopodia.




8

8 *Vorticella* (60 μm), phylum Protista, class Ciliata. Each has conspicuous cilia (forming a spiral) that beat food particles into the gullet, a meganucleus, a micronucleus, a contractile vacuole and stalks with a central contractile filament.




9

9 *Volvox* (1 mm), a colonial green alga, class Chlorophyceae. Each individual resembles a single *Chlamydomonas*. There are daughter colonies inside the main colonies.



10

10 *Spirogyra* (40 μm and 100 μm diameter), a filamentous green alga, class Chlorophyceae. The spirally arranged chloroplasts are typical.



11

11 *Paramecium* (250 μm), phylum Protista, class Ciliata. Note the cilia, the gullet, food vacuoles, the meganucleus and two contractile vacuoles (one with collecting channels).



12

12 *Micrasterias* (300 μm), a desmid, class Chlorophyceae. This unicellular green alga has a pair of cellulose cases forming two 'semi-cells'. It looks similar to a diatom, but the pigments are different and the cases are of different materials.

13 Flowers of *Rubus* (blackberry) with a butterfly (*Lycaena phloea*, order Lepidoptera) sipping nectar. The plant is an angiosperm (family Rosaceae) with open flowers and many stamens.



13

14 *Tortula muralis*, a moss, superclass Bryophyta. The sporophytes are tall with slender capsules at their tips from which spores are released. The gametophyte forms a cushion on rocks and walls.



14



15



16



17



18



19



20

19

15 A branch of a larch tree *Larix*, subclass Gymnospermidae. The cones are old female cones which have opened, showing seeds.

16 *Polytrichum*, a moss, superclass Bryophyta (seen from above). Note the shoot tips of the gametophytes with male reproductive organs.

17 A branch of birch *Betula*, subclass Angiospermidae. This is wind-pollinated and the flowers form catkins of two sexes. Male flowers form the catkin that hangs down and the small female catkin is upright.

18 *Lamium album* (white deadnettle). The flowers are complex in shape with stamens arched under the upper petal; when bees alight on the lower lip they are brushed with pollen.

19 *Marchantia*, a liverwort, superclass Bryophyta. The small cups on the upper surface are gemmae, in which small vegetative propagules develop. The umbrella-like structures carry male reproductive organs.

20 *Equisetum*, a horsetail, class Sphenopsida. The central thick shoot bears reproductive structures (sporangia). The leaves are arranged in whorls at the nodes.

21 *Lycopodium clavatum*, a club moss, class Lycopsidea. These leafy shoots arise from a rhizome and carry sporangia in the axils of the leaves near the tips.



21



22



23

22 *Ganoderma applanata* (bracket fungus). This is the fruiting body of a basidiomycete that grows in dead and dying trees.

23 Flowers of the honeysuckle *Lonicera periclymenum*; these are pollinated by insects. They are of complex shape with nectaries at the base of a long tube.

24 *Peziza coccineus* (a cup fungus). This is the fruiting body of an ascomycete.

25 Knapweed (*Centaurea*) visited by a bumble bee *Bombus hortorum*. This 'flower' is a collection of similar florets, each with stamens and stigmas. The bee belongs to the order Hymenoptera and is already dusted with pollen.

26 *Morchella vulgaris* (a morel). This is an edible fruiting body of an ascomycete.

27 Flowers of *Heracleum* (hogweed) with a visiting beetle (*Strangalia quadrifasciata*, order Coleoptera). This plant belongs to the order Umbelliferae and has characteristic flower heads.

28 *Armillaria mellea* (honey fungus). These are fruiting bodies of a basidiomycete fungus that lives on rotting wood.

29 *Amanita muscaria* (fly agaric). This is the fruiting body of a basidiomycete fungus and it is very poisonous.

30 *Phallus impudicus* (stinkhorn), the fruiting body of a basidiomycete. The spores are dispersed by flies that are attracted by the smell.

31 A tipulid (daddy-long-legs), order Diptera. Note the haltere behind the left wing.



24



25



26



27



28



29



30



31



34



35



37



32



36



33

32 Puss moth caterpillar (*Cercora vinula*, order Lepidoptera). There are three pairs of short thoracic legs and paired abdominal 'pro-legs'.

33 Burnet moths (*Zygaena filipendula*, order Lepidoptera) copulating. The female emerged from the pupal case (right). This poisonous moth flies by day.

34 *Coriza hyoscyami*, a bug (order Hemiptera). Note the forewings.

35 *Peripatus*, a primitive uniramian from Australia. The soft body looks velvety because of the many tracheal openings.

36 *Ophrys apifera*, a bee orchid, mimicking a female bee; male bees try to copulate with the flowers.

37 A black carabid beetle (order Coleoptera) with a small worker ant (order Hymenoptera).

38 *Meconema*, a bush cricket (order Orthoptera). Note the membrane forewings.



38



39

39 *Rana temporaria*, a common frog, class Amphibia.



40

40 *Lacerta vivipara*, the sand lizard, class Reptilia.

41 Life on a rocky shore. A picture showing typical animals and plants of British rocky shores.



41



42



43

42 *Chaetodon* (butterfly fish), a teleost that lives among coral reefs, seen in front of a coral colony, with tentacles expanded.

43 *Triturus vulgaris*, the smooth newt, class Amphibia.

44 Life on a sandy shore. A picture showing typical animals of British sandy shores and of the sea above.



44



45



46



47



48




49



50



51



52

45 *Laminaria* (tangle, oar-weed), a brown seaweed that grows on the lower shore and in shallow seas. There is a narrow stipe and a broad, flat lamina.

46 *Fucus serratus* (serrated wrack), a brown seaweed that lives on the lower part of the middle shore. It lacks bladders.

47 *Ascophyllum nodosum* (knotted wrack), a brown seaweed, with *Polysiphonia* (a red seaweed) growing on it. Both live on the middle shore. Note the large air-bladders and narrow thallus of the wrack.

48 *Fucus vesiculosus* (bladder wrack), a brown seaweed that lives on the middle shore. Note the pairs of air-bladders and the flat thallus.

49 Smooth periwinkles *Littorina littoralis* on fronds of *Fucus vesiculosus*. This mollusc species has many colour forms including bright yellow.


50 *Pelvetia canaliculata* (channelled wrack), a brown seaweed that lives on the upper shore. The channels are on the underside of the thallus; the swollen tips contain reproductive organs.

51 Barnacles (*Chthamalus stellatus*) are crustaceans with calcareous shell plates attached to rocks.

52 Rough periwinkles *Littorina saxatilis* (phylum Mollusca, class Gastropoda) retreat to crevices when the tide goes out.

53 Common limpets *Patella vulgata* (phylum Mollusca, class Gastropoda).


54 Dog whelk *Nucella lapillus* with its egg cases attached to the rocks beside it. This carnivorous gastropod lives on the middle shore.



53



54




55A


55A Electron micrograph of a whole rat liver cell showing nucleus (N), nucleolus (Nu), nuclear membrane (NM), mitochondria (M), endoplasmic reticulum (ER) (ribosomes are present on the endoplasmic reticulum but are barely visible at this low magnification), cell membranes (CM), bile canaliculus (BC) (part of the bile duct system) and glycogen granules (G). The Golgi apparatus is not visible in the section. ($\times 9\,500$)

56B Light micrographs of sections of the intestines of mice injected with thymidine: (a) 8 hours after injection; (b) 36 hours after injection. TN, tagged nuclei; V, villus. ($\times 70$)


57C Electron micrograph of cell membranes (CM) of intestinal cells showing the three-layered membrane. DOL, dense outer layer; CIL, clear inner layer; DIL, dense inner layer; IS, intercellular space. ($\times 160\,000$)




56Ba



56Bb



57C



58D

58D Electron micrograph of freeze-etched cell membranes (CM) from two neighbouring cells in the pancreas. The inner face of the membrane (arrowed) contains many intramembraneous particles. ($\times 250\,000$)

59E Electron micrograph of the junctional complex between adjacent lateral cell membranes (LCM) of rat intestinal cells. In the region of the tight junction (TJ) the apposed membranes often appear fused though there is a small gap here. IJ, intermediate junction; D, desmosome; ACM, apical cell membrane; Mv, microvillus. ($\times 105\,000$)




59E




60F Electron micrograph of a gap junction (G) in an ovarian granulosa cell. ($\times 30\,000$)

61G Electron micrograph of the boundary between adjacent corn (*Zea*) cells showing plasmodesmata (arrowed) and mitochondria (M). ($\times 18\,000$)




61G




62H

62H Electron micrograph showing phagocytosis by a polymorphonuclear leucocyte (L). Note the evagination (E) around a micro-organism (M) and another micro-organism enclosed within the cell (*). ($\times 40\,000$)




63I

63I Electron micrograph of a liver cell showing nucleus (N), nucleolus (Nu), nuclear membranes (NM), nuclear pore (NP), rough endoplasmic reticulum with ribosomes (RER). G, Golgi apparatus; M, mitochondria; DB, dense bodies. ($\times 27\,500$)

**64J**


64J Electron micrograph of a liver cell showing nuclear membranes (NM) with ribosomes on the cytoplasmic side, nuclear pores (NP), extensive rough endoplasmic reticulum (RER) and mitochondria (M). ($\times 45\,000$)




65K

65K Electron micrograph of a freeze-etched preparation from onion. Note the pores (*arrowed*) seen in both transverse and cross-section. N, nucleus; C, cytoplasm; G, Golgi apparatus; M, mitochondria. ($\times 25\ 000$)


66L Electron micrograph of a cell from the human liver. The left-hand side of the field is packed with the twisted cisternae of the smooth endoplasmic reticulum (SER); at the bottom right is some rough endoplasmic reticulum (RER); also on the right are mitochondria (M). ($\times 32\ 000$)






67M

67M Electron micrograph of the Golgi apparatus. ($\times 47\,000$)



68N



69O




70P


68N Electron micrograph of a liver cell showing dense bodies believed to be lysosomes (L), reaction product (RP) indicating the site of acid phosphatase activity and a mitochondrion (M). ($\times 25\,000$)

69O Electron micrograph of two peroxisomes from a cell of the grass plant *Avena* showing the single external membrane (M) and the core of densely staining, crystal-like material (C). ($\times 108\,000$)

70P Electron micrograph showing mitochondria (M) from a liver cell with transverse or longitudinal cristae (C). Note the rough endoplasmic reticulum (RER) surrounding the mitochondria. ($\times 88\,000$)



71Qa




71Qb

71Q (a) Electron micrograph of isolated cristae from a mitochondrion swollen in a hypotonic solution and negatively stained with phosphotungstate. ($\times 150\,000$)

(b) An enlargement showing the elementary (F_1) particles (*arrowed*). ($\times 650\,000$)


72R Electron micrograph of an immature plant cell showing proplastids (P). Note also the cell wall (CW) and vacuole (V). ($\times 2\,000$)



72R



73S




74T


73S Electron micrograph of a chloroplast showing stacks of grana (G) within the stroma (S). A mitochondrion (M) is on the left. ($\times 25\ 000$)

74T Electron micrograph of a pair of centrioles in a Chinese hamster cell, at right angles to each other. TSC, transversely sectioned centriole showing nine triplets of subunits skewed towards the centre: LSC, longitudinally sectioned centriole. ($\times 37\ 000$)

75U Electron micrograph of cilia in transverse section showing the '9 + 2' arrangement of tubules (T). Note the short 'arms' on the A tubules of each doublet, and the membrane (M) surrounding the cilia. ($\times 180\ 000$)



75U




76V

76V Electron micrograph of mitochondria in a rat liver cell fixed with potassium permanganate. ($\times 86\,000$)



77W

77W Electron micrograph of mitochondria in a rat liver cell fixed conventionally with glutaraldehyde followed by osmium tetroxide. ($\times 86\,000$)




78a

78 (a) A model showing the three-dimensional structure of ribonuclease. The polypeptide chain is shown as white with hydrogen bonds as red bars; negatively charged, amino acid sidechains are shown as blue spheres, positively charged sidechains as red spheres and sulphur atoms concerned in disulphide bonding as prominent yellow spheres. Tyrosine, tryptophan and other hydrophobic amino acids are shown as purple spheres.

Note that the overall shape of the molecule is a V, with the active site cleft between the two arms of the V (this cleft runs vertically from the top of the molecule). You should also be able to see that the structure is held together by hydrogen bonds in both of the common secondary structures, the β -pleated sheet and α -helix. The most obvious region of an α -helix is close to the N-terminus (amino terminus; see (b)) and forms one face of the cleft. This helix runs from top left to bottom right, and you should be able to see the hydrogen bonds parallel to the direction of the helix. The bottom right-hand edge of the molecule contains a large area of β -pleated sheet, consisting of three chains held together by hydrogen bonds perpendicular to the direction of the chain.

(b) The folding of the polypeptide backbone in ribonuclease. The orientation of the molecule is the same as in (a). Disulphide bridges are shown as solid black circles and active site histidines as aromatic ring structures.




78b




79

79 *Drosophila melanogaster*. This small fruitfly (about 2.5 mm long) is widely used in developmental biology and genetics because of its ready availability and rapid, if complicated, life cycle.



80





81

80 Light micrograph of the human pancreas. Insulin-containing cells have been labelled with anti-insulin antibodies. The antibodies have then been shown up by a histochemical technique. This reveals the islet (endocrine) regions (Is) as darkly stained areas surrounded by exocrine cells (ExC). ($\times 65$)


81 Low-power electron micrograph of the human pancreas showing endocrine cells at the edge of an islet. Glucagon-secreting cells (A cells) and insulin-secreting cells (B cells) can be recognized from their distinctive ultrastructure. ExC, exocrine cells; Is, islet. ($\times 3\,300$)

82 High-power electron micrograph of a steroid-secreting cell from the *zona fasciculata* region of the rat adrenal cortex. Note the lipid droplets (L) and abundant smooth endoplasmic reticulum (SER) which is often closely associated with the droplets. ($\times 66\,000$) The inset is a low-power view of the cell showing the distinctive mitochondria (M). The structural appearance of these varies in the different regions of the gland but they all possess the characteristic tubular cristae. N, nucleus. ($\times 23\,000$)





83





84

83 Endocrine cells in the anterior pituitary of the laboratory mouse. Particular cell types can be identified on the basis of the size and density of the secretory granules and by using immunohistochemistry techniques. This micrograph shows secreting cells of the following types: prolactin (PRL), adrenocorticotropic hormone (ACTH), luteinizing hormone (LH) and growth hormone (GH). The endocrine cells are either adjacent to a capillary (Cap) or an intercellular space (IS) that connects with the capillary. ($\times 5\,400$)


84 The ultrastructure of a synapse. This synapse is between two neurons from the cerebellum of a young rat. Note the density of synaptic vesicles in the presynaptic ending, the thickening of both presynaptic and postsynaptic membranes and the synaptic cleft. ($\times 160\,000$)

85 The neurohaemal region in the rat posterior pituitary. Neurosecretory nerve endings (NE) can be seen next to an endothelial cell (E) that forms part of a capillary (Cap). Neurosecretory axons often have swollen regions which are used as storage depots for the neurosecretion. One of these swellings (S) can be seen in the centre of the picture. Note the large number of neurosecretory granules (NG). ($\times 10\,000$)





86



87



88



89a



89b

86 Electron micrograph of a capillary from the brain (hippocampus) of a rat, shown in cross-section. The large dark object in the centre is a single, red blood cell. ($\times 10\,000$)

87 Blood and lymph capillaries in the mesentery of a rat. The lymphatics are the large, dark vessels; the blood vessels are the very much smaller vessels. Note the blind ends of the lymphatics. The preparation was specially treated to make the lymphatics stand out—usually they are colourless and cannot be seen so well. ($\times 55$)


88 Photomicrograph of rat muscle. Carbon has been injected intravenously to show up the vascular pattern. The main arteries and veins can be seen running parallel to one another (veins are the larger vessels). Note the artery–artery shunt (*red dashed line*), vein–vein shunt (*blue dashed line*), and also that this is a very different type of capillary bed from that found in the mesentery. ($\times 15$ approx.)

89 Valves of a lymphatic vessel. (a) Partial contraction of a collecting lymphatic in rat mesentery. (b) 3 seconds later the vessel is in a relaxed state. Note the capillary alongside. ($\times 160$)

90 A photographic reconstruction of a capillary from arteriole to venule in the cat mesentery. Direct recordings of the mean blood pressure (expressed as kPa) were taken at points A–F. The capillaries range from 7.5 to 9 μm in diameter.

91 Interstitial volume can increase to a gross extent when lymphatic drainage is prevented. In this case, lymphatics from the scrotum are blocked by millions of parasites—filarial worms—and the condition, for obvious reasons, is known as elephantiasis.


92 The membrane between the blood and the cavity of Bowman's capsule. The lumen of the capillary (Cap) lies to the left; the cavity of Bowman's capsule (BC) lies to the right. The endothelium has numerous fenestrations (F). The slit-pores between the foot processes are closed by a thin membrane (M). The principal structure separating blood from glomerular filtrate is the basement membrane (BM).



90




91



92




93a




93b

93 (a) Scanning electron micrograph of three gill filaments (F) of *Mytilus edulis* showing the frontal cilia (C). Note that the latero-frontal cirri (LFC) extend across the interfilamentary spaces to form a fine-mesh filter. ($\times 1\ 100$)

(b) High-power view of the latero-frontal cirri in (a). Each latero-frontal cirrus consists of a double row of 20–25 pairs of cilia (C), one of each pair being on either side of the main axis of the cirrus. The effect is to form a mesh-work between the cirri and also between adjacent filaments. ($\times 8\ 000$)




94



95

94 Electron micrograph of sieve-plate pores in the phloem of *Aristolochia brasiliensis*. The pores are occluded with P-protein but relatively free from callose. ($\times 64\,000$)

95 Electron micrograph showing a sieve tube and sieve plate from *Nicotiana tabaccum* (tobacco) in longitudinal section. The sieve-plate pores appear open. P, sieve-plate pore; CC, companion cell; PI, plastid; PP, P-protein. ($\times 5\,000$)




96




97

96 Electron micrograph of a sieve tube of *Nicotiana* in transverse section, PP, P-protein. ($\times 1\ 800$)


97 Transverse section of the stem of *Tilia* (lime) showing the penetration of an aphid stylet (AS) into a single sieve-tube element (S). ($\times 800$)



98



99




100


98 Mammalian kidney showing glomerulus surrounded by the glomerular capsule (*arrowed*). The capsule is made up of squamous epithelium. ($\times 300$)

99 Squamous epithelium (*arrowed*) lining secondary lamellae of the gills of a teleost fish. ($\times 400$)


100 Endothelium lining the lumen of an artery. ($\times 125$)



101



102




103


101 Cuboidal epithelium lining the wall of a thyroid gland follicle. ($\times 400$)

102 Cuboidal epithelium is also found in the submucosa of the small intestine where it forms Brunner's glands. ($\times 400$)

103 Cuboidal epithelium from kidney collecting-tubules. ($\times 600$)



104




105

104 Columnar epithelium on a villus from the small intestine of a rat. A histochemical reaction has stained mucus-producing cells a greenish colour. ($\times 600$)

105 Ciliated, columnar epithelium lining the trachea of a mammal; the epithelium has a pseudo-stratified appearance. ($\times 400$)


106 Lightly keratinized epithelium. ($\times 400$)




106



107a



107b




108


107 (a) Low-power view of heavily keratinized epithelium from the palm of the hand. ($\times 160$)

(b) High-power view of the central area of (a). ($\times 400$)

108 Transitional epithelium lining the inside wall of bladder. ($\times 400$)



109



110a



110b


109 Dense connective tissue of the type encountered in the supporting tissues of the skin. ($\times 400$)

110 (a) Low-power view of loose connective tissue from the bladder of a frog. ($\times 200$)

(b) High-power view of the central area of (a). ($\times 500$)



111



112



113

111 Hyaline cartilage from mammalian trachea. ($\times 80$)

112 A transverse section of compact bone. ($\times 170$)

113 Mammalian smooth muscle in LS. ($\times 500$)


114 Three arrangements of tissue from a mammal. At the top is smooth muscle in TS; beneath this is an area containing a large parasympathetic nerve cell; at the bottom is smooth muscle in LS. ($\times 500$)



114



115a




115b

115 (a) Skeletal muscle fibres in LS. ($\times 750$)

(b) Skeletal muscle fibres in TS. Note the peripheral location of the nuclei. ($\times 350$)

116 Cardiac muscle in LS. Note the branching nature of the fibres. ($\times 450$)



116



117



118



119


117 A leaf of *Pinus* (Scots pine), a xerophyte, in TS.

118 A leaf of *Prunus* (cherry), a mesophyte, in TS.

119 A leaf of *Nymphaea* (waterlily), a hydrophyte, in TS.



120



120a



120b




120c




120d


120 Photographs of the external genitalia of mature female ferrets. The top photograph shows the position in which the four lower photographs were taken. (a)–(d) are four groups of ferrets kept under different conditions.




121



122



123



124

121 Dorsal view of the heart of a pig. The diameter of the pulmonary arterial wall appears larger than usual, partly as a result of the angle of the heart and partly because the pulmonary artery had been slashed. The ventricles are foreshortened because of the angle at which the picture was taken. LA, left atrium; RA, right atrium; LV, left ventricle; RV, right ventricle.


122 Pig's heart, showing the result of slicing across the apex of the two ventricles.

123 The pig's heart opened to display the right ventricular cavity.


124 The left ventricular chamber of the pig's heart.



125a



125b



125c



125d

125 Cucumber roots in TS, taken 10 μm behind the root tip. Seedlings were germinated in (a) water, (b) 0.1 mg l^{-1} 2,4-D, (c) 1 mg l^{-1} 2,4-D, and (d) 10 mg l^{-1} 2,4-D. In (d), note that only part of the whole section is shown. All photographs are at the same magnification. ($\times 400$)

index to plates

a

A cells 81 28
ACTH-secreting cells 83 30
adrenal cortex 82 29
Amanita (fly agaric) 29 6
Amoeba 7 3
amphibian 39 8; 43 9
Anabaena 1 3
angiosperms 13 4; 17, 18 5;
23, 25, 27 6; 36 7
anterior pituitary 83 30
aphid stylet 97 36
Armillaria (honey fungus) 28 6
artery-artery shunt 88 32
ascomycete 24, 26 6
Ascophyllum 47 10

b

Bacillariophyceae 2 3
barnacles (*Chthamalus*) 51 11
basidiomycete 22, 28, 29, 30 6
Betula (birch) 17 5
B cells 81 28
bivalves 44 9
blood capillary (rat brain) 86 32
blood capillary (rat mesentery) 87,
89 32
blood capillary (cat mesentery) 90 33
Bombus (bumble bee) 25 6
Bowman's capsule: membrane 92 33
Bryophyta 14 4; 16, 19 5

c

carabid beetle 37 7
Centaurea (knapweed) 25 6
Ceratium 5 3
Chaetodon (butterfly fish) 42 9
Chlamydomonas 6 3
Chlorophyceae 4, 6 3; 9, 10, 12 4
chloroplast 73S 23
Chrysophyta 2 3
ciliate 8, 11 4
cnidarian 41 8; 42, 44 9
Coleoptera 27 6; 37 7
Coriza (bug) 34 7
Crustaceans 41 8; 44 9; 51 11
Cyanophyceae 1, 4 3

d

daddy-long-legs 31 7
Dinophyceae 5 3
Diptera 30 6; 31 7; 79 27
Drosophila melanogaster 79 27

e

echinoderms 44 9
elephantiasis 91 33
endocrine cells (anterior pituitary)
83 30
endocrine cells (pancreas) 80, 81 28
Equisetum (horsetail) 20 5
Euglena 3 3
exocrine cells (pancreas) 80, 81 28

f

fish 42, 44 9
Fucus serratus 46 10
Fucus vesiculosus 48, 49 10

g

Ganoderma (bracket fungus) 22 6
gastropods 49 10; 52, 53, 54 11
GH-secreting cells 83 30
gill filaments (*Mytilus edulis*) 93 34
gymnosperm 15 5

h

Hemiptera 34 7; 97 36
Heracleum (hogweed) 27 6
Hymenoptera 25 6; 37 7

i

islets 80, 81 28

l

Lacerta (sand lizard) 40 8
Laminaria 45 10
Lamium (deadnettle) 18 5
latero-frontal cirri of gill filament
93 34
Larix (conifer) 15 5
LH-secreting cells 83 30
Lepidoptera 13 4; 32, 33 7
lipid droplet 82 29
Littorina littoralis (smooth
periwinkle) 49 10
Littorina saxatilis (rough
periwinkle) 52 11
Lonicera (honeysuckle) 23 6
Lycaena (butterfly) 13 4
Lycopodium (club moss) 21 5
Lycopsida 21 5
lymph capillary 87 32
lymphatic valves 89 32

m

Marchantia (liverwort) 19 5
Meconema (bush cricket) 38 7
mitochondrion 61G 14; 65K
18; 70P 21; 71Q 22
Micrasterias 12 4
molluscs 41 8; 44 9; 49 10;
52, 53, 54 11
Morchella (morel) 26 6

n

Navicula 2 3
neurosecretory granule 85 31
neurosecretory nerve 85 31
neurohaemal region 85 31
Nucella (dog whelk) 54 11

o

Ophrys (bee orchid) 36 7
Orthoptera 38 7

p

pancreas **58D** 13; **80, 81** 28
Paramecium **11** 4
Patella (common limpet) **53** 11
Pelvetia **50** 10
Peripatus **35** 7
Peziza (cup fungus) **24** 6
Phallus (stinkhorn) **30** 6
Phaeophyta **45, 46, 47, 48, 50** 10
phloem **94, 95** 35; **96, 97** 36
plasmodesmata **61G** 14
Polytrichum (moss) **16** 5
postsynaptic membrane **84** 30
posterior pituitary **85** 31
P-protein **94, 95** 35; **96** 36
puss moth caterpillar **32** 7
PRL-secreting cells **83** 30
presynaptic membrane **84** 30
protistans **7** 3; **8, 11** 4
Pyrrophyta **5** 3

r

Rana (common frog) **39** 8
reptile **40** 8
ribonuclease structure **78** 26
Rhodophyta **47** 10
rocky shore organisms **41** 8
Rubus (blackberry) **13** 4

S

sandy shore organisms **44** 9
Sarcodina **7** 3
seaweeds **41** 8
secretory granules **83** 30
sieve plate **94, 95** 35
sieve tube **95** 35; **96, 97** 36
single-celled blue-green **4** 3
smooth endoplasmic reticulum **82** 29
Spirogyra **10** 4
sphenopsid **20** 5
steroid-secreting cell **82** 29
Strangalia (beetle) **27** 6
synapse **84** 30
synaptic vesicles **84** 30
synaptic cleft **84** 30

t

teleost **42, 44** 9
Tortula (moss) **14** 4
Triturus (smooth newt) **43** 9

U

uniramian **35** 7

V

vein-vein shunt **88** 32
Volvox **9** 4
Vorticella **8** 4

W

worker ant **37** 7

Z

Zygnema **4** 3
Zygaena (burnet moth) **33** 7

acknowledgements

Grateful acknowledgement is made to the following for illustrations used in this book.

1-12 Dr J. W. G. Lund; 13-21, 23, 25, 27, 31-34, 36-40, 45-48, 50, 51 and 54 J. E. Bebbington (Field Studies Council); 22, 24, 26, 28-30 Dr I. Ridge (Open University); 35, 42, 49, 52, 53 and 79 Dr M. E. Varley (Open University); 43 Dr T. Halliday (Open University); 55A, 57C, 59E, 61G, 63I, 64J, 72R and 73S Dr M. G. Stewart, Dr P. Mullins and J. Broadbent (Open University); 56B Professor C. P. Leblond (McGill University); 58D Professor L. Orci (University of Geneva); 60F E. A. Anderson (Harvard Medical School); 62H Professor D. Bainton (University of California); 65K Professor D. Branton, (in *Cells and Organisms*, A. B. Novikoff, 1970, Holt, Rinehart & Winston); 66L Churchill Livingstone (Medical Division of Longman Group Ltd) for Dr P. Toner (in *Cell Structure*, Toner and Carr, 2nd edn, 1971); 67M C. F. Leedale (in *An Atlas of Biological Ultrastructure*, J. B. Dodge, 1968, Arnold); 69O A. B. Novikoff and E. Holtzman, *Cells and Organelles*, 1976, Holt, Rinehart & Winston; 70P Professor B. R. Brinkley (University of Texas); 71Q H. Fernandez-Moran (in *Cell Biology*, De Robertis *et al.*, 6th edn, 1976, W. B. Saunders); 74T Professor B. R. Brinkley and E. Stubblefield (in *Cells and Organisms*, A. B. Novikoff, 1976, Holt, Rinehart & Winston); 75U B. A. Afzelius, *Journal of Ultrastructure Research*, 37, 1971; 76V and 77W Dr M. G. Stewart and J. Broadbent (Open University); 80 and 81 Dr A. Clarke and D. Jerome (John Radcliffe Hospital, Oxford); 82 Dr G. Bullock (Ciba-Geigy Pharmaceuticals); 83 Professor R. L. Holmes (University of Leeds); 84 Dr R. Beale (Open University); 85 Dr J. F. Morris (University of Oxford), *Journal of Endocrinology*, 68, 1976; 86-90 Professor B. W. Zweifach; 91 U.S. Armed Forces Institute of Pathology; 92 Professor D. Moffatt; 93 Professor G. Owen (University of Aberystwyth); 94 H. D. Behnke; 95 and 96 J. Cronshaw; 97 M. H. Zimmerman; 98-116 and 125 Open University Histological Laboratory; 117-119 Gene Cox.

List of Units

The Diversity of Organisms

Unit 1	Marine Organisms
Unit 2	From Sea to Land: Plants and Arthropods
Unit 3	From Sea to Land: Vertebrates

Cell Biology

Unit 4	Cell Structure
Unit 5	Macromolecules and Membranes
Unit 6	Enzymes: Specificity and Catalytic Power
Unit 7	Enzymes: Regulation and Control
Unit 8	Metabolism and its Control
Units 9 and 10	Membrane Transport and Energy Exchange

Development

Unit 11	Development: The Component Processes
Units 12 and 13	Cellular Differentiation
Unit 14	Pattern Specification and Morphogenesis
Unit 15	Chicken or Egg?

Animal Physiology

Unit 16	Communication: Nerves and Hormones
Unit 17	Blood Sugar Regulation
Unit 18	Control Mechanisms in Reproduction
Unit 19	Circulatory Systems
Unit 20	Respiratory Mechanisms
Unit 21	Respiratory Gases and Their Transport
Units 22 and 23	Osmoregulation and Excretion
Units 24 and 25	Nutrition, Feeding Mechanisms and Digestion

Plant Physiology


Unit 26	Plant Water Relations
Unit 27	Ion Movement and Phloem Transport
Unit 28	Plants and Energy
Unit 29	Plant Cells: Growth and Differentiation
Units 30 and 31	Morphogenesis in Flowering Plants



The Open University Press



S202 Picture Book



0 335 16042 5